

Image Article

Leser - Trelat Sign in Adenocarcinoma of Gallbladder

Samir Jha*

Bassett Medical Center, Cooperstown, New York, USA



Figure 1: Multiple variably shaped seborrheic keratoses as seen on the back.



Figure 2: Keratoses involving neck and the upper chest.

Clinical Image

A 93-year-old, frail lady presented to our hospital with multiple seborrheic keratoses over her entire back and chest. She reported a sudden and rapid appearance of the skin lesions before histopathological confirmation of gall bladder adenocarcinoma. At this visit, she reported the progression of seborrheic keratoses with wide-spreading of lesion over back (Figure 1) and seborrheic keratoses lesions on her chest (Figure 2), correlating with metastatic involvement of hepatic capsule.

*Corresponding author: Samir Jha, Bassett Medical Center, Cooperstown, New York, USA; E-mail: samir.jha@bassett.org

Citation: Samir Jha (2019) Leser - Trelat Sign in Adenocarcinoma of Gallbladder. *Int J Clin Med Imaging* 6: 636. doi:10.4172/2376-0249.1000636

Copyright: © 2019 Samir Jha. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.