

Case Blog

Title: Superficial Basal Cell Carcinoma

Alex H Wong¹, Alexander KC Leung^{2*} and Benjamin Barankin³

¹Department of Family Medicine, University of Calgary, Canada

²Department of Pediatrics, University of Calgary; Pediatric Consultant, Alberta Children's Hospital, Canada

³Department of Dermatology, Toronto Dermatology Centre, Canada



A 62-year-old Caucasian woman presented with a slowly growing, asymptomatic, red plaque on her back that was first noted a year ago. Her family physician treated her with a topical antifungal and subsequently a topical steroid without success. Basal cell carcinoma (BCC) is generally defined as a locally invasive, slowly spreading tumor that rarely metastasizes, arising in the epidermis or hair follicles and in which, in particular, the peripheral cells usually simulate the basal cells of the epidermis. BCCs are a heterogeneous group of tumors that based on their clinical morphology can be classified most commonly into the following subtypes: nodular BCC, superficial BCC, sclerosing BCC, infiltrative BCC, micronodular BCC, and pigmented BCC. Superficial BCC is the second most common subtype, accounting for 9 to 17.5% of all cases. Unlike other BCC subtypes, it tends to have an earlier age of onset and a female predominance. It is the least aggressive subtype. Typically, the lesion presents as a solitary, well-defined, slowly growing, erythematous or pinkish, scaly patch or plaque that has a thin pearly border. The size varies from a few millimeters to several centimeters in diameter. The condition is usually asymptomatic. Occasionally, the lesion may be pigmented or ulcerated. Sites of predilection include the trunk, shoulders, and extremities. Diagnosis is often delayed as it is often mistaken for a dermatitis or tinea infection. For lesions occurring at a low-risk anatomic site (e.g., trunk and extremities) with distinct clinical borders and sizes less than 2 centimeters in diameter, treatment options include surgical excision, electrodesiccation with curettage, liquid nitrogen cryotherapy, topical immunotherapy (e.g. imiquimod), topical 5-fluorouracil, and photodynamic therapy. Patients should have regular follow-up at least yearly following diagnosis to monitor for local recurrence or the development of new BCCs or other skin cancers at other sites.

*Corresponding author: Alexander K.C. Leung, MBBS, FRCPC, FRCP(UK and Ireland), FRCPCH, FAAP, Clinical Professor of Pediatrics, the University of Calgary, #200, 233 – 16th Avenue NW Calgary, Alberta, T2M 0H5, Canada, Tel: (403) 230-3322; E-mail: aleung@ucalgary.ca

Copyright: © 2015 Wong et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.